

86-A186 014

ON THE FEYNMAN-KAC'S FORMULA AND ITS APPLICATIONS TO
FILTERING THEORY. (U) NORTH CAROLINA UNIV AT CHAPEL
HILL CENTER FOR STOCHASTIC PROC R L KARANDIKAR
OCT 86 TR-161 AFOSR-TR-87-1096


1/1

UNCLASSIFIED

F/G 12/3

NL





ROCOPY RESOLUTION TEST CHART

DTIC FILE COPY

UNCLASSIFIED

AD-A186 014

①

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE


1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA			
4. PERFORMING ORGANIZATION REPORT NUMBER -		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TM- 87-1096	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL If applicable:	7a. NAME OF MONITORING ORGANIZATION AFOSR NM	
6c. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 0324A, Chapel Hill, NC 27514		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL If applicable: NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
11. TITLE (Include Security Classification) ON THE FEYNMAN-KAC'S FORMULA AND ITS APPLICATIONS TO FILTERING THEORY		TASK NO. A5	WORK UNIT NO.
12. PERSONAL AUTHOR(S) R.L. Karandikar			
13a. TYPE OF REPORT Technical	13b. TIME COVERED 9/30/85- 9/30/86 FROM -XXXXXXXXXXXXX	14. DATE OF REPORT (Yr., Mo., Day) Oct. 86	15. PAGE COUNT 21
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Keywords:	
FIELD GROUP SUB GR			
XXXXXXXXXXXXXXXXXX			
19. ABSTRACT Let (X_t) be a Markov process, not assumed to be time homogeneous. It is well known that $\hat{X}_t = (t, X_t)$ is a time homogeneous Markov process. Let A be its generator. The Feynman-Kac's formula for X_t takes the following form if the equation (1.1) $Av + cv = 0$ admits a solution v , then v has the representation, for $s < t$, (1.2) $v(s, X_s) = E \left[v(t, X_t) \exp \left(\int_s^t c(u, X_u) du \right) \mathcal{G}(X_s) \right]$. We prove this under general conditions on (X_t) .			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Peggy Rawitch		22b. TELEPHONE NUMBER (Include Area Code) 919-962-2307	22c. OFFICE SYMBOL AFOSR/NM

DTIC
ELECTE
OCT 07 1987
S E D

CENTER FOR STOCHASTIC PROCESSES

AFSR-TR. 87-1096

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



ON THE FEYNMAN-KAC'S FORMULA AND ITS APPLICATIONS TO FILTERING THEORY

by

Rajeeva L. Karandikar

Technical Report No. 161

October 1986

ON THE FEYNMAN-KAC's. FORMULA AND ITS APPLICATIONS TO FILTERING THEORY

RAJEEVA L. KARANDIKAR

Center for Stochastic Processes
University of North Carolina
and

Indian Statistical Institute
7, S.J.S. Sansanwal Marg, New Delhi



Technical Report No. 161 (UNC)

Technical Report No. 8408 (ISI)

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Research partially supported by the Air Force Office of Scientific
Research Contract #F49620 85 C 0144 and by the Indian Statistical
Institute.

ON THE FEYNMAN-KAC'S. FORMULA AND ITS APPLICATIONS TO FILTERING THEORY

Rajeeva L. Karandikar
Indian Statistical Institute
7, S.J.S. Sansanwal Marg, New Delhi
and
Center for Stochastic Processes
University of North Carolina

1. Introduction : Let (X_t) be a Markov process, not assumed to be time homogeneous. It is well known that $\hat{X}_t = (t, X_t)$ is a time homogeneous Markov process. Let A be its generator. The Feynman-Kac's formula for X_t takes the following form if the equation

$$(1.1) \quad Av + cv = 0$$

admits a solution v , then v has the representation, for $s < t$

$$(1.2) \quad v(s, X_s) = E \left[v(t, X_t) \exp \left(\int_s^t c(u, X_u) du \right) \middle| \sigma(X_s) \right].$$

We prove this under general conditions on (X_t) .

Then we come to the question of existence of solution to (1.1). We show that under some regularity conditions on (X_t) , (1.1) has a solution for a rich class of boundary conditions. This implies that the 'dual' equation to (1.1) admits a unique solution. The 'dual' equation is an equation for measures on the state space of (X_t) and its unique solution is the distribution of X_t under an absolutely continuous change of the underlying probability measure by a multiplicative functional.

These results on the measure valued equations significantly extend results given in [3] on the conditional distributions for the nonlinear filtering problem (in the white noise approach).

2. Let (S, \underline{S}) be a measurable space. Let (X_t) be an (S, \underline{S}) valued Markov process on a probability space $(\Omega, \underline{A}, \pi)$ with transition probability function P , i.e.

$$\{\omega : X_t(\omega) \in B\} \in \underline{A}$$

and

$$(2.1) \quad E_{\pi} \left[1_B(X_t) | \underline{F}_s^X \right] = P(s, X_s, t-s, B) \quad \text{a.s. } \pi$$

for all $0 \leq s \leq t < \infty$, $B \in \underline{S}$. Here, the function $P(s, x, t, B)$ on $\{0 \leq s < \infty, t \geq 0, x \in S, B \in \underline{S}\}$ is assumed to satisfy the following conditions.

(2.2) For $s \geq 0, t \geq 0, x \in S$; $P(s, x, t, \cdot)$ is a countably additive probability measure on (S, \underline{S}) .

(2.3) For $s \geq 0, x \in S, B \in \underline{S}$; $P(s, x, 0, B) = 1_B(x)$.

(2.4) For $t \geq 0, B \in \underline{S}$; $(s, x) \rightarrow P(s, x, t, B)$ is a $B([0, \infty)) \otimes \underline{S}$ measurable function ($\underline{B}(E)$ denotes the Borel σ -field of a topological space E and \otimes denotes the product of σ -fields).

(2.5) For $s \geq 0, u \geq 0, t \geq 0, x \in S, B \in \underline{S}$; we have

$$\int_S P(s+t, z, u, B) P(s, x, t, dz) = P(s, x, t+u, B)$$

Throughout, \underline{F}_t^X denotes the smallest σ -field with respect to which the family $\{X_u : 0 \leq u \leq t\}$ is measurable. We also assume that

(2.6) the process (X_t) is \underline{F}_t^X -progressively measurable, i.e. for all $t_0 < \infty$, the mapping $(t, \omega) \rightarrow X_t(\omega)$ from $[0, t_0] \times \Omega \rightarrow S$ is $\underline{B}([0, t_0]) \otimes \underline{F}_{t_0}^X$ measurable.

: 3 :

Let $\hat{S} = [0, \infty) \times S$, $\hat{S}_t = \mathbb{B}([0, \infty)) \otimes S_t$ and \underline{J} be the class of bounded real valued \hat{S}_t measurable functions on \hat{S} .

Definition : A sequence $\{f_k\} \subseteq \underline{J}$ is said to converge weakly to $f \in \underline{J}$, written as $w\text{-}\lim_{k \rightarrow \infty} f_k = f$, if $f_k(x)$ is uniformly bounded and for each $x \in S$, $f_k(x)$ converges to $f(x)$.

For $f \in \underline{J}$, $t \geq 0$, let $T_t f : S \rightarrow \mathbb{R}$ be defined by

$$(2.7) \quad (T_t f)(s, x) = \int f(s+t, z) P(s, x, t, dz), \quad (s, x) \in \hat{S}.$$

Using the properties of P , it can be checked that $T_t f \in \underline{J}$ and that for $u \geq 0$, $t \geq 0$,

$$(2.8) \quad T_u [T_t f] = T_{t+u} f, \quad f \in \underline{J}.$$

Thus $\{T_t : t \geq 0\}$ is a semigroup of operators (from \underline{J} into itself).

Remark : It is well known and easy to check that $\hat{X}_t = (t, X_t)$ is a Markov process with stationary transition probability function \hat{P} given by

$$\hat{P}(t, (s, x), B) = P(s, x, t, B^{(s+t)}) , \quad B \in \hat{\mathcal{S}}$$

where B^u denotes the u -section of $B \subseteq S$. The semigroup $\{T_t : t \geq 0\}$ defined above is the usual semigroup associated with the transition function P (as in [2], section 2.1).

We will now recall the definition and some properties of the weak generator A of $\{T_t : t \geq 0\}$. Let \underline{J}_0 be given by

$$\underline{J}_0 = \{f \in \underline{J} ; \lim_{t \downarrow 0} T_t f = f\}$$

Definition : Let \underline{D}_A be the class of $f \in \underline{J}$ for which the

$$(2.9) \quad \lim_{t \downarrow 0} \frac{T_t f - f}{t} = g$$

exists and belongs to \underline{J}_0 and for $f \in \underline{D}_A$, define $Af = g$, where g is given by (2.9).

The following properties are easy to prove. We will only state them here. For a proof see chapter 1 in [2].

$$(2.10) \quad T_t(\underline{D}_A) \subseteq \underline{D}_A \quad \text{and for } f \in \underline{D}_A, \quad A(T_t f) = T_t Af$$

$$(2.11) \quad \text{For } f \in \underline{J}_0, \quad t \mapsto (T_t f)(s, x) \text{ is a right continuous function for all } (s, x) \in \hat{S}.$$

$$(2.12) \quad \text{For } f \in \underline{D}_A, \text{ we have, for all } (s, x) \in \hat{S}, \quad t \geq 0$$

$$(T_t f)(s, x) = f(s, x) + \int_0^t (T_u Af)(s, x) du$$

(2.13) Given $f \in \underline{J}_0$ there exists a sequence $\{f_k\} \subseteq \underline{D}_A$ such that

$$\lim_{k \rightarrow \infty} f_k = f.$$

In (2.13) above, f_k can be taken to be

$$f_k(s, x) = \int_0^\infty k e^{-kt} (T_t f)(s, x) dx.$$

The property (2.12) has the following important consequence.

Proposition 1 : For $f \in \underline{D}_A$, M_t given by

$$(2.14) \quad M_t(\omega) = f(t, X_t(\omega)) - \int_0^t (Af)(u, X_u(\omega)) du$$

is a martingale with respect to the σ -fields \underline{F}_t^X .

Proof : The progressive measurability of (X_t) implies the \underline{F}_t^X measurability of M_t . Since $f, Af \in \underline{J}$, they are bounded and hence M_t is itself bounded for each t . Now (2.1) implies

$$(2.15) \quad \begin{aligned} E_\pi [f(t, X_t) | \underline{F}_s^X] &= \int f(t, z) P(s, X_s, t-s, dz) \\ &= (T_{t-s} f)(s, X_s) \end{aligned}$$

for $s \leq t$. Similarly for $s \leq u$, we have

$$(2.16) \quad E_\pi [(Af)(u, X_u) | \underline{F}_s^X] = (T_{u-s} Af)(s, X_s).$$

Using (2.11), (2.12), (2.15) and (2.16), it can be checked that

$$E_\pi [M_t - M_s | \underline{F}_s^X] = 0.$$

□

We now turn our attention to the Feynman-Kac's formula. Our next result is a step in this direction.

Let $g : [0, t_0] \times S \rightarrow \mathbb{R}$ be a $\mathbb{B}([0, t_0]) \times \mathcal{S}$ measurable function such that

$$(2.17) \quad E_{\pi} \left[\int_0^{t_0} |g(u, X_u)| du \right] < \infty$$

and for a positive integrable function $a : [0, t_0] \rightarrow \mathbb{R}$,

$$(2.18) \quad g(u, x) \leq a(u) \quad \text{for all } x \in S, u \in [0, t_0].$$

Fix $0 \leq s \leq t_0$ and let

$$(2.19) \quad B_t(\omega) = \exp\left(\int_s^t g(u, X_u(\omega)) du\right).$$

Then we have

Theorem 2 : Let $f \in \underline{D}_A$ and g satisfy (2.17), (2.18). Then

$$(2.20) \quad Z_t = f(t, X_t) \cdot B_t - \int_s^t [(Af)(u, X_u) + g(u, X_u)] \cdot B_u du$$

is an \mathbb{F}_t^X martingale for $t \geq s$ (where B is given by (2.19)).

Proof : It is easy to see that Z_t is \mathbb{F}_t^X measurable. The condition (2.18) implies that B_t is bounded. Since f, Af are also bounded the condition (2.17) gives the integrability of Z_t . To prove the martingale property, suffices to prove that for $s \leq r \leq t$, $C \in \mathbb{F}_r^X$,

$$(2.20) \quad E_{\pi} [(Z_t - Z_r) \cdot 1_C] = 0.$$

Let $f_1(t, \omega) = f(t_0, X_{t_0}(\omega)) - \int_t^{t_0} (Af)(u, X_u(\omega)) du$. Then by Proposition 1, it follows that for $0 \leq t \leq t_0$

: 7 :

$$(2.21) \quad E_{\pi} \left[f_1(t, \cdot) | \mathcal{F}_t^X \right] = f(t, X_t)$$

and hence

$$(2.22) \quad E_{\pi} \left[1_C \cdot (Z_t - Z_r) \right] = E_{\pi} 1_C \cdot \{ f_1(t, \cdot) B_t - f_1(r, \cdot) B_r - \int_r^t \{ (Af + gf)(u, X_u) du \} \}.$$

Now for each ω , $f_1(t, \omega)$, $B_t(\omega)$ are absolutely continuous functions and hence

$$\begin{aligned} f_1(t, \omega) B_t(\omega) - f_1(r, \omega) B_r(\omega) &= \int_r^t \frac{d}{du} [f_1(u, \omega) B_u(\omega)] du \\ &= \int_r^t \{ f_1(u, \omega) \cdot g(u, X_u(\omega)) B_u(\omega) \\ &\quad + (Af)(u, X_u(\omega)) \cdot B_u(\omega) \} du. \end{aligned}$$

Thus

$$\begin{aligned} (2.23) \quad E_{\pi} [f_1(t, \cdot) B_t - f_1(r, \cdot) B_r] &= E 1_C \int_r^t f_1(u, \cdot) g(u, X_u) B_u du \\ &\quad + E 1_C \int_r^t (Af)(u, \cdot) B_u du \\ &= E 1_C \int_r^t (Af + gf)(u, X_u) du \end{aligned}$$

using (2.21) once again. Now (2.22) and (2.23) give the required equality

$$E [1_C (Z_t - Z_r)] = 0.$$

Remark : It can be verified that

$$Z_t = M_t B_t - \int_s^t M_u dB_u$$

where M is given by (2.14). Hence if M were right continuous,

it would follow from the "integration by parts formula for martingale"

(See [5]) that (Z_t, \mathbb{F}_t^X) is a martingale. However, in general M_t need not be right continuous and hence we have given a direct proof.

The following is the Feynman-Kac's formula for a time inhomogeneous Markov process.

Theorem 3 : Let $0 < t_0 < \infty$ be fixed. Let $c : [0, t_0] \times S \rightarrow \mathbb{R}$ and $g : S \rightarrow \mathbb{R}$ be bounded measurable functions. Suppose that $v \in \mathcal{D}_A$ is a solution to

$$(2.24) \quad [Av + cv](u, x) 1_{\{u < t_0\}} = 0$$

and

$$(2.25) \quad v(t_0, x) = g_0(x).$$

Then v admits a representation, for $s < t_0$

$$(2.26) \quad v(s, X_s) = E_{\pi} \left[g_0(X_{t_0}) \exp \left(\int_s^{t_0} c(u, X_u) du \right) \middle| \mathbb{F}_s^X \right] \quad \text{a.e. } \pi.$$

Proof : Fix $s < t_0$. Take $f = v$ and $g = c$ in Theorem 2 to obtain that $(Z_t, \mathbb{F}_t^X)_{s \leq t \leq t_0}$ is a martingale, where

$$(2.27) \quad Z_t = v(t, X_t) \exp \left(\int_s^t v(u, X_u) du \right).$$

Here we have used the fact that v satisfies (2.24) so that the second term appearing in the expression for Z_t is zero. Thus

$$E_{\pi} [Z_{t_0} | \mathbb{F}_s^X] = Z_s \quad \text{a.s. } \pi$$

This is same as (2.26) since $v(t_0, x) = g_0(x)$.

□

3. In this section we consider the question as to under what conditions in $c, g, (X_t)$ does the problem (2.24), (2.25) admit a solution. Of course, if the solution exists, it has to satisfy (2.26) and this gives a clue as to what conditions one should put on $c, g, (X_t)$.

Suppose that S is a topological space, \mathcal{S} is its Borel σ field. Let \mathcal{X} be the space of all right continuous mappings \underline{X} from $[0, \infty)$ into S . We will denote by \underline{X}_t the value of \underline{X} at t . Let $\mathcal{F}_t^S = \sigma(\underline{X}_u : s \leq u \leq t)$. We assume that

$$(3.1) \quad \text{for all } \omega, \underline{X}_s(\omega) \in \mathcal{X}$$

and that for all $(s, x) \in \hat{S}$, there exists a probability measure $P_{s,x}$ on $(\mathcal{X}, \mathcal{F}_\infty^S)$ such that for $0 \leq t_0 \leq t_1 \leq \dots \leq t_k, y_i \in S, A_1, A_2, \dots, A_k \in \mathcal{S}$; $k \geq 1$, we have

$$(3.2) \quad P_{t_0, y}(\underline{X}_{t_i} \in A_i : 1 \leq i \leq k) = \int \dots \int \prod_{i=1}^k 1_{A_i}(y_i) P(t_{i-1}, y_{i-1}, t_i - t_{i-1}, dy_i).$$

Remark The main thrust of this assumption is that $P_{s,x}$ is realized on \mathcal{X} . The relation (2.1) and (3.2) imply that for $\{t_i\}, \{A_i\}$ as in (3.2), we have

$$(3.3) \quad E_\pi \left[\prod_{i=1}^k 1_{A_i}(\underline{X}_{t_i}) \middle| \mathcal{F}_{t_0}^X \right] = P_{t_0, \underline{X}_{t_0}}(\underline{X}_{t_i} \in A_i : 1 \leq i \leq k) \text{ a.s. } \pi$$

and hence by standard arguments, we have for $B \in \mathcal{A}_{\infty}^{t_0}$,

$$(3.4) \quad \pi(\underline{X} \in B | \mathcal{F}_{t_0}^X) = P_{t_0, \underline{X}_{t_0}}(B) \quad \text{a.s. } \pi.$$

Similarly, it can be proved that for $s < t, B \in \mathcal{A}_{\infty}^t, x \in S$,

$$(3.5) \quad P_{s,x}(B | \mathcal{A}_t^S) = P_{t, \underline{X}_t}(B) \quad \text{a.s. } P_{s,x}.$$

We are now in a position to prove a 'converse' to the Feynman-Kac's formula.

Theorem 4 : Let $0 < t_0 < \infty$ be fixed. Let $c : [0, t_0] \times S \rightarrow \mathbb{R}$ be a bounded continuous function. Let $f \in \underline{D}_A$. Let $v : \hat{S} \rightarrow \mathbb{R}$ be defined by

$$(3.6) \quad \begin{aligned} v(s, x) &= E_{P_{s, x}} \left[f(t_0, X_{t_0}) \exp \left(\int_s^{t_0} c(u, X_u) du \right) \right], \quad s < t_0 \\ &= f(s, x), \quad s \geq t_0. \end{aligned}$$

Then $v \in \underline{D}_A$ and $Av = f_1$ where

$$(3.7) \quad \begin{aligned} f_1(s, x) &= -c(s, x)v(s, x), \quad s < t_0 \\ &= (Af)(s, x), \quad s \geq t_0. \end{aligned}$$

Proof Since $v(s, x) = f(s, x)$ for $s \geq t_0$, we have

$$(T_t v)(s, x) = (T_t f)(s, x)$$

for $s \geq t_0$, $x \in S$, $t \geq 0$. Hence for $s \geq t_0$, $x \in S$.

$$(3.8) \quad \lim_{t \downarrow 0} \frac{(T_t v)(s, x) - v(s, x)}{t} = (Af)(s, x) = f_1(s, x).$$

For $x \in X$, $s \leq t_0$ let us define

$$(3.9) \quad C_s(x) = \exp \left(\int_s^{t_0} c(u, X_u) du \right).$$

Then for $s \leq t_0$, we have

$$v(s, x) = E_{P_{s, x}} \left[f(t_0, X_{t_0}) C_s(x) \right].$$

: 11 :

For $s < t_0$, $s + t < t_0$, we thus have

$$\begin{aligned}
 (3.10) \quad (T_t v)(s, x) &= \int v(s+t, z) P(s, x, t, dz) \\
 &= E_{P_{s, x}} \left[v(s+t, \underline{X}_{s+t}) \right] \\
 &= E_{P_{s, x}} \left[E_{P_{s+t, \underline{X}_{s+t}}} (f(t_0, \underline{X}_{t_0}) C_{s+t}(\underline{X})) \right] \\
 &= E_{P_{s, x}} \left[f(t_0, \underline{X}_{t_0}) C_{s+t}(\underline{X}) \right].
 \end{aligned}$$

by (3.5). Hence, for $s < t_0$, $x \in S$, $s + t < t_0$ we have

$$(3.11) \quad \frac{(T_t v)(s, x) - v(s, x)}{t} = E_{P_{s, x}} \left[f(t_0, \underline{X}_{t_0}) \cdot \frac{C_{s+t}(\underline{X}) - C_s(\underline{X})}{t} \right].$$

For all $\underline{X} \in \mathcal{X}$, we have from (3.9)

$$(3.12) \quad \lim_{t \rightarrow 0} \frac{C_{s+t}(\underline{X}) - C_s(\underline{X})}{t} = -c(s, \underline{X}_s) \cdot C_s(\underline{X}).$$

Further

$$(3.13) \quad \left| \frac{C_{s+t}(\underline{X}) - C_s(\underline{X})}{t} \right| = \left| -c(t, \underline{X}_t) \cdot C_t(\underline{X}) \right|$$

$$\leq K$$

where K depends only on t_0 and the upper bound of $|c|$. The dominated convergence theorem gives that for $s < t_0$

$$\begin{aligned}
 (3.14) \quad \lim_{t \rightarrow 0} \frac{(T_t v)(s, x) - v(s, x)}{t} &= E_{P_{s, x}} \left[f(t_0, \underline{X}_{t_0}) \{-c(s, \underline{X}_s) C_s(\underline{X})\} \right] \\
 &= -c(s, x) v(s, x) \\
 &= f_1(s, x)
 \end{aligned}$$

as $P_{s,x}(X_s = x) = 1$. Also, (3.13) implies that the left hand expression in (3.11) is uniformly bounded (in s, x, t). Thus we have

$$(3.15) \quad \lim_{t \downarrow 0} \frac{(T_t v)(s, x) - v(s, x)}{t} = f_1(s, x).$$

Remains to prove that $f_1 \in J_0$. This will prove that $v \in D_A$ and that $Av = f_1$. If $s \geq t_0$, $f_1(s, x) = (Af)(s, x)$ and hence for $s \geq t_0$, $t \geq 0$, $(T_t f_1)(s, x) = (T_t Af)(s, x)$. Since $Af \in J_0$, this gives

$$(3.16) \quad \lim_{t \downarrow 0} (T_t f_1)(s, x) = f_1(s, x) \quad \text{for } s \geq t_0, x \in S.$$

For $s < t_0$, we have

$$\begin{aligned} T_t f_1(s, x) - f_1(s, x) &= -E_{P_{s,x}} \left[c(s+t, X_{s+t}) v(s+t, X_{s+t}) - c(s, x) v(s, x) \right] \\ &= -E_{P_{s,x}} \left[v(s+t, X_{s+t}) \{c(s+t, X_{s+t}) - c(s, x)\} \right] \\ &\quad - c(s, x) E_{P_{s,x}} [v(s+t, X_{s+t}) - v(s, x)]. \end{aligned}$$

Now as $t \downarrow 0$, $c(s+t, X_{s+t}) \rightarrow c(s, X_s) = c(s, x)$ a.e. $P_{s,x}$, as c is continuous and X_s is right continuous. Hence by the dominated convergence theorem,

$$\lim_{t \downarrow 0} E_{P_{s,x}} [v(s+t, X_{s+t}) \{c(s+t, X_{s+t}) - c(s, x)\}] = 0.$$

The relation (3.14) implies that

$$-c(s, x) E_{P_{s,x}} [v(s+t, X_{s+t}) - v(s, x)] = -c(s, x) [(T_t v)(s, x) - v(s, x)]$$

$\rightarrow 0$

as $t \downarrow 0$. These observations give

: 13 :

$$(3.17) \quad \lim_{t \rightarrow 0} (T_t f_1)(s, x) = f_1(s, x) \quad \text{for } s < t_0, x \in S.$$

Now (3.16), (3.17) and the fact that $T_t f_1$ is uniformly bounded yield

$$w\text{-}\lim_{t \rightarrow 0} (T_t f_1) = f_1.$$

□

Remark : Under the conditions assumed in this section and Theorem 4, the equations (2.24), (2.25) for $g_0(x) = f(t_0, x)$ have a unique solution v on $[0, t_0] \times S$ which is given by (3.6). To see this, let v' be any solution. Apply Theorem 3 to the process $\{X_t : t \geq s\}$ on the probability space $(X, \underline{A}_\infty^s, P_{s,x})$ to obtain, for $s < t_0$,

$$v'(s, X_s) = E_{P_{s,x}} \left[f(t_0, X_{t_0}) \mid \underline{A}_s^s \right] \quad \text{a.s. } P_{s,x}.$$

Since under $P_{s,x}$ any set in \underline{A}_s^s has measure zero or one, the conditional expectation appearing above is the unconditional expectation and thus equals $v(s, x)$. Also $X_s = x$ a.s. $P_{s,x}$. Hence we have

$$v'(s, X_s) = v(s, x) \quad \text{a.s. } P_{s,x}.$$

These observations imply

$$v'(s, x) = v(s, x).$$

4. We now consider an equation dual to (2.24), namely

$$\frac{d}{dt} K_t = A^* K_t + g(t, \cdot) K_t$$

where $\{K_t\} \subseteq \underline{M}(S)$ - the class of finite signed measures on (S, \underline{S}) . The equation (4.1) is purely formal and is to be interpreted as

$$(4.2) \quad \langle f(t, \cdot), K_t \rangle = \langle f(0, \cdot), K_0 \rangle + \int_0^t \langle Af(u, \cdot), K_u \rangle du + \int_0^t \langle g(u, \cdot) f(u, \cdot), K_u \rangle du$$

for $f \in \underline{D}_A$. Here, $\langle \theta, \mu \rangle$ denotes $\int \theta d\mu$ for $\mu \in \underline{M}(S)$ and a function $\theta: S \rightarrow \mathbb{R}$. Thus $\langle f(t, \cdot), \mu \rangle = \int f(t, x) d\mu(x)$ for $f \in \underline{J}$. We will show that this equation with boundary condition

$$(4.3) \quad K_0 = \Pi \circ X_0^{-1}$$

admits a unique solution which is given by

$$(4.4) \quad K_t(B) = E_\pi \left[1_B(X_t) \exp \left(\int_0^t g(u, X_u) du \right) \right], \quad B \in \underline{S}.$$

The uniqueness will be proved in the class of $\{K_t\}$ satisfying

$$(4.5) \quad \{K_t\} \subseteq \underline{M}(S), \quad t \mapsto K_t(B) \text{ is a Borel measurable function}$$

for all $B \in \underline{S}$ and $K_t \ll \Pi \circ X_t^{-1}$ with

$$\left| \frac{dK_t}{d\Pi \circ X_t^{-1}} \right| \leq M$$

for all t , for a fixed constant M .

We continue to assume that the conditions imposed on (X_t) in Section 3, are valid. We further assume that S is a complete separable metric space. We begin with a Lemma.

Lemma 5 : Let $0 < t < \infty$ be fixed. Let $\mu \in \underline{M}(S)$ be such that

$$(4.6) \quad \langle f(t, \cdot), \mu \rangle = 0 \quad \forall f \in \underline{D}_A.$$

Then $\mu \equiv 0$.

Proof : Let \underline{E} be the class of $f \in \underline{J}$ for which (4.6) holds. Easy to see that if $f_k \in \underline{E}$, $w\text{-}\lim_{k \rightarrow \infty} f_k = f$, then $f \in \underline{E}$. Hence by (2.13), $\underline{J}_0 \subseteq \underline{E}$.

For $f \in C_b(\hat{S})$, (i.e. $f : \hat{S} \rightarrow \mathbb{R}$ is bounded continuous), we have

$$(T_t f)(s, x) = E_{P_{s, x}} f(s+t, X_{s+t}) \rightarrow f(s, x) \text{ as } t \downarrow 0,$$

since X_u is right continuous. Thus $C_b(\hat{S}) \subseteq \underline{J}_0 \subseteq \underline{E}$.

Given $f_0 \in C_b(S)$, taking $f(s, x) = f_0(x)$, we have $f \in C_b(\hat{S}) \subseteq \underline{E}$ and hence

$$(4.7) \quad \langle f_0, \mu \rangle = 0.$$

The validity of (4.7) for all $f_0 \in C_b(S)$ implies $\mu = 0$ because \underline{S} - the Borel σ field - is also the smallest σ field with respect to which $C_b(S)$ is measurable.

□

We are now in a position to prove the assertions made at the beginning of this section. This result may be considered as a dual Feynman-Kac's formula.

Theorem 6 : Suppose that g satisfies (2.17) and (2.18). Then the equation (4.2) with boundary condition (4.3) admits a unique solution in the class of $\{K_t\}$ satisfying (4.5). The unique solution is given by (4.4).

Proof : First we will prove that $\{K_t\}$ defined by (4.4) satisfies (4.2). Let $\{K_t\}$ be defined by (4.4). Easy to see that (4.3) and (4.5) are satisfied.

Taking $s = 0$ in Theorem 2, it follows from the martingale property of Z_t that $E_\pi Z_t = E_\pi Z_0$. Here, Z_t is given by (2.20) where in turn B_t is given by (2.19), with $s = 0$. Noting that with these notations,

$$\langle \theta, K_t \rangle = E_\pi \theta(X_t) B_t$$

we conclude from the relation $E_\pi Z_t = E_\pi Z_0$ that

$$\langle f(t, \cdot), K_t \rangle - \int_0^t \langle (Af + \sigma f)(u, \cdot), K_u \rangle du = \langle f(0, \cdot), K_0 \rangle.$$

Hence $\{K_t\}$ satisfies (4.2).

To prove the uniqueness part, we will prove the following. Suppose $\{K_t\}$ satisfies (4.2), (4.5) and $K_0 \equiv 0$. Then $K_t \equiv 0$, $t \geq 0$.

For this fix $t_0 < \infty$ and $f \in \underline{D}_A$. Let ν be the measure defined on $S' = [0, t_0] \times S$ by

$$(4.8) \quad \nu(B) = E_\pi \int_0^{t_0} 1_B(u, X_u) du, \quad B \in \underline{S}' = \underline{S}(S').$$

Then note that (2.17) implies $\int_{S'} |g| d\nu < \infty$. Hence if $g_k : S' \rightarrow \mathbb{R}$ is defined by

$$(4.9) \quad g_k(s, x) = g(s, x) 1_{\{|g(s, x)| \leq k\}}$$

then we have

$$(4.10) \quad \int_{S'} |g_k - g| d\nu \rightarrow 0 \quad \text{as} \quad k \rightarrow \infty.$$

For each k , g_k is bounded by k . By Lusin's theorem (see [1], p. 187) we can get $c_{k,i} \in C_b(S')$, bounded by k , such that

$$(4.11) \quad c_{k,i} \rightarrow g_k \quad \text{a.e. } \nu \quad \text{as } i \rightarrow \infty.$$

Hence

$$(4.12) \quad \lim_{i \rightarrow \infty} \int |c_{k,i} - g_k| dv = 0.$$

Let $v_{k,i}$ be given by (3.6) for $c = c_{k,i}$. Then $A v_{k,i} = -c_{k,i} v_{k,i}$ on $[0, t_0) \times S$ by Theorem 4. Using (4.2) for $v_{k,i}$ and recalling that $K_0 = 0$, we have

$$(4.13) \quad \begin{aligned} \langle f(t_0, \cdot), K_{t_0} \rangle &= \langle v_{k,i}(t_0, \cdot), K_{t_0} \rangle \\ &= \int_0^{t_0} \langle (A v_{k,i} + g v_{k,i})(u, \cdot), K_u \rangle du \\ &= \int_0^{t_0} \langle (g - c_{k,i}) v_{k,i}(u, \cdot), K_u \rangle du. \end{aligned}$$

Thus

$$(4.14) \quad \begin{aligned} |\langle f(t_0, \cdot), K_{t_0} \rangle| &\leq M \int_0^{t_0} \langle |(g - c_{k,i}) v_{k,i}|(u, \cdot), \pi \circ X_u^{-1} \rangle du \\ &= M E_\pi \int_0^{t_0} |(g - c_{k,i}) v_{k,i}|(u, X_u) du. \\ &= M \int |g - c_{k,i}| \cdot |v_{k,i}| dv. \end{aligned}$$

As $i \rightarrow \infty$, $v_{k,i}$ converges pointwise to v_k and is bounded by k , where v_k is given by (3.6) for $c = g_k$. This and (4.11), (4.12), (4.14) imply

$$|\langle f(t_0, \cdot), K_{t_0} \rangle| \leq M \int |g - g_k| \cdot |v_k| dv.$$

Since (4.9) implies $c_k(u, x) \leq a(u)$, it follows that

$$|v_k| \leq M_1 \cdot \exp\left(\int_0^{t_0} a(u) du\right) = M_2$$

where $|f| \leq M_1$. Hence

$$|\langle f(t_0, \cdot), K_{t_0} \rangle| \leq M \cdot M_2 \int |g - g_K| dv.$$

This and (4.10) imply $\langle f(t_0, \cdot), K_{t_0} \rangle = 0$. Since $f \in \mathcal{D}_A$ is arbitrary, Lemma 5 gives $K_{t_0} = 0$. This completes the proof. \square

We will briefly consider the equation for normalized measures

$$(4.15) \quad N_t(B) = \frac{K_t(B)}{K_t(S)}, \quad B \in \mathcal{S}$$

where K_t is given by (4.4). It is easy to see, using (4.2) that $\{N_t\}$ satisfies.

$$(4.16) \quad \langle f(t, \cdot), N_t \rangle = \langle f(0, \cdot), N_0 \rangle + \int_0^t \langle (Af + gf)(u, \cdot), N_u \rangle du - \int_0^t \langle f(u, \cdot), N_u \rangle \langle g(u, \cdot), N_u \rangle du$$

We will now prove that $\{N_t\}$ is the unique solution to this equation.

Theorem 7 : The equation (4.16) with boundary condition $N_0 = \mathbb{W} \circ X_0^{-1}$ admits a unique solution in the class of $\{N_t\}$ satisfying (4.5). The solution is given by (4.15).

Proof : We need to prove uniqueness of the solution. Let N_t^i be any other solution, i.e. satisfying (4.5), (4.16) and $N_0^i = \mathbb{W} \circ X_0^{-1}$. Then it can be checked that $N_t^i(S) = 1$ for all $t \geq 0$. Further, if K_t^i is defined by

$$(4.17) \quad K_t^i(B) = N_t^i(B) \cdot \exp\left(\int_0^t \langle g(u, \cdot), N_u^i \rangle du\right),$$

then K_t^i is a solution to (4.2) and that it satisfies (4.3), (4.5).

Hence by Theorem 6, $K_t^i = K_t$. This and the observation that

$$N_t^i(B) = \frac{K_t^i(B)}{K_t^i(S)}$$

give us the required equality, namely $N_t^i = N_t$. \square

5. We will now give applications of the results in the previous sections to filtering theory.

We refer the reader to [4] for a detailed discussion and background on the white noise approach to filtering theory.

We assume that the signal process (X_t) is a Markov process satisfying the conditions imposed in the previous sections.

Let \underline{K} be a separable Hilbert space. Let $h : [0, T] \times S \rightarrow \underline{K}$ be a measurable function such that

$$(5.1) \quad E_{\pi} \left[\int_0^T \|h_u(X_u)\|_{\underline{K}}^2 du \right] < \infty$$

Let $H = L^2([0, T], \underline{K})$ and let $\xi: \Omega \rightarrow H$ be defined by

$$(\xi(\omega))_u = h_u(X_u(\omega)), \quad 0 \leq u \leq T.$$

Consider the model

$$y = \xi + e$$

where $e = (e_t)$ is \underline{K} -valued white noise independent of (X_t) . Here y is the observation process and y, ξ, e are realised on a Quasi cylinder probability space $(E, \underline{E}, \alpha)$ (See [4] section 6). We now state the Bayes formula. For the relevant definitions and proof, see [4].

Theorem 8 : For $g : S \rightarrow \mathbb{R}$ bounded, measurable,

$$(5.2) \quad E_{\alpha}(g(X_t) | y_u : u \leq t) = \int_S g(x) dF_t(y)(x)$$

where

$$(5.3) \quad F_t(y)(B) = E_{\pi} \left[1_B(X_t) \exp \left(\int_0^t (h_u(X_u), y_u)_{\underline{K}} du - \frac{1}{2} \int_0^t \|h_u(X_u)\|_{\underline{K}}^2 du \right) \right]$$

and

$$(5.4) \quad F_t(y)(B) = \frac{\Gamma_t(y)(B)}{\Gamma_t(y)(S)}$$

for $0 \leq t \leq T$, $y \in H$, $B \in \underline{S}$.

$\Gamma_t(y)$, $F_t(y)$ are known as unnormalized and normalized conditional distribution of X_t given $\{y_u : u \leq t\}$, respectively.

The following is an immediate consequence of Theorems 6,7. Let $\varepsilon_y(t,x) = (h_t(x), y_t)_{\underline{K}} - \frac{1}{2} \|h_t(x)\|_{\underline{K}}^2$, $(t,x) \in [0,T] \times S$, $y \in H$.

Theorem 9 : Let h satisfy (5.1). (i) For all $y \in H$, $\Gamma_t(y)$ is the unique solution to the equation

$$(5.5) \quad \langle f(t, \cdot), \Gamma_t(y) \rangle = \langle f(0, \cdot), \Gamma_0(y) \rangle + \int_0^t \langle (Af + \varepsilon_y f)(u, \cdot), \Gamma_u(y) \rangle du,$$

$f \in \underline{D}_A$

with the condition $\Gamma_0(y) = \pi \circ X_0^{-1}$ in the class of $\{K_t\}$ satisfying (4.5).

(ii) For all $y \in H$, $F_t(y)$ is the unique solution to the equation

$$(5.6) \quad \langle f(t, \cdot), F_t(y) \rangle = \langle f(0, \cdot), F_0(y) \rangle + \int_0^t \langle (Af + \varepsilon_y f)(u, \cdot), F_u(y) \rangle du \\ - \int_0^t \langle f(u, \cdot), F_u(y) \rangle \langle \varepsilon_y(u, \cdot), F_u(y) \rangle du, \quad f \in \underline{D}_A$$

with the initial condition $F_0(y) = \pi \circ X_0^{-1}$ in the class of $\{K_t\}$ satisfying (4.5).

Proof : Since

$$|\varepsilon_y(t,x)| \leq \|h_t(x)\|_{\underline{K}}^2 + \frac{1}{2} \|y_t\|_{\underline{K}}^2$$

and

$$\varepsilon_y(t,x) \leq \frac{1}{2} \|y_t\|_{\underline{K}}^2,$$

it follows that for all $y \in H$, ε_y satisfies (2.17) and (2.18). Thus (i) follows from Theorem 6 and (ii) from Theorem 7.

Remark : Theorem 9 was proved in [3] under the much stronger condition

$$(5.7) \quad ||h_t(x)|| \leq a_t \quad \text{with} \quad \int_0^T a_t^2 dt < \infty .$$

The equations (5.5) and (5.6) are analogues of the Zakai and Fujisaki-Kallianpur-Kunita equations. In [3], $r_t(y)$ and $F_t(y)$ were also characterized as unique solutions to another type of equations (equations (3.4) and (3.11) in [3]) under the condition (5.7). With a little bit of work, it can be shown that (5.7) can be replaced by (5.1) in these results as well.

References

1. Ash, R. : Real Analysis and Probability, Academic Press, New York (1972).
2. Dynkin, E.B. : Markov processes, Vol. 1, Springer-Verlag, Berlin (1965)
3. Kallianpur, G. and Karandikar, R.L. : Measure valued equations for the optimum filter in finitely additive nonlinear filtering theory. Z.Wahrscheinlichkeitstheorie verw. Geb., 66, 1-17 (1984).
4. Kallianpur, G. and Karandikar, R.L. : White noise calculus and nonlinear filtering theory. The Annals of Probability 13, 1033-1107 (1985).
5. Stroock, D.W. and Varadhan, S.R.S. : Multidimensional diffusion processes, Springer-Verlag, Berlin (1979).

END

DATE
FILMED

DEC.

1987